ELSEVIER

Environmental Pollution 153 (2008) I–VI

ENVIRONMENTAL POLLUTION

ww.elsevier.com/locate/envpol

Contents of Volume 153

Number 1 2008

SPECIAL ISSUE SECTION: Chelating Agents in the Environment

Guest editorial

1 **Chelating agents and the environment**
B. Nowack

3 **The use of chelating agents in the remediation of metal-contaminated soils: A review**
D. Leštan, C.-l. Luo, X.-d. Li

The use of synthetic chelants for soil washing and enhanced phytoextraction by plants has been well studied for the remediation of metal-contaminated soils in the last two decades.

14 **Pressure-assisted chelation extraction of lead from contaminated soil**
P.K.A. Hong, X. Cai, Z. Cha

Chelant extraction of metals from contaminated soil is enhanced by rapid, successive cycles of air compression and immediate decompression wherein pressure and number of cycles are key parameters.

22 **Electrokinetic remediation of concrete: Effect of chelating agents**
K. Popov, I. Glazkova, V. Yachmenev, A. Nikolayev

EK remediation in combination with chelating agents has a good potential for contaminated concrete surfaces.

29 **Promising role of plant hormones in translocation of lead in *Sesbania drummondii* shoots**
M. Israr, S.V. Sahi

Plant hormones (IAA and NAA) enhance the accumulation of lead in shoots of Sesbania drummondii.

37 **Formation of aerobic granules in the presence of a synthetic chelating agent**
Y.V. Nancharaiah, H.M. Joshi, T.V. Krishna Mohan, V.P. Venugopalan, S.V. Narasimhan

Synthetic chelating agent enhances aerobic microbial granulation.

44 **Influence of variable chemical conditions on EDTA-enhanced transport of metal ions in mildly acidic groundwater**
D.B. Kent, J.A. Davis, J.L. Joye, G.P. Curtis

Metal-ion adsorption strongly influences transport of metal–EDTA complexes in groundwater.

53 **Glyphosate behavior at soil and mineral–water interfaces**
R.C. Pessagno, R.M. Torres Sánchez, M. dos Santos Afonso

Adsorption isotherms, surface coverage and zeta potential curves of glyphosate in aqueous suspensions of montmorillonite and three Argentine soils were measured as a function of PMG concentration and pH.

Regular papers

60 **Variation in PAH inputs and microbial community in surface sediments of Hamilton Harbour: Implications to remediation and monitoring**
G.F. Slater, B.R. Cowie, N. Harper, I.G. Droppo

Variations in PAH inputs to harbour sediments have implications to implementation and monitoring of mitigation/remediation efforts.

71 **Toxaphene in minke whales (*Balaenoptera acutorostrata*) from the North Atlantic**
B. Gouteux, D.C.G. Muir, S. Backus, E.W. Born, R. Dietz, T. Haug, T. Metcalfe, C. Metcalfe, N. Øien

High levels of toxaphene were found in different sub-populations of minke whales from North Atlantic waters.

84 **Organochlorine residue levels in livers of birds of prey from Spain: Inter-species comparison in relation with diet and migratory patterns**
B. van Drooge, R. Mateo, Í. Vives, I. Cardiel, R. Guitart

The contribution of bird biomass in the diet is a determining factor for the accumulation of organochlorines in raptors.

92 **Distribution of organochlorine pesticides (OCPs) in conifer needles in the southeast Tibetan Plateau**
R. Yang, T. Yao, B. Xu, G. Jiang, X. Zheng

Conifer needles in southeastern Tibet indicate several organochlorine pesticides are transported from the Indian continent by the southern Asian monsoon.

doi:10.1016/S0269-7491(08)00221-2

101 Plasma vitellogenin in Morelet's crocodiles from contaminated habitats in northern Belize
T.R. Rainwater, K.W. Selcer, L.M. Nespoli, A.G. Finger, D.A. Ray, S.G. Platt, P.N. Smith, L.D. Densmore, T.A. Anderson, S.T. McMurry

Wild crocodiles living in habitats polluted with organochlorine pesticides did not exhibit contaminant-induced vitellogenin induction in blood plasma.

110 Bioavailable DDT residues in sediments: Laboratory assessment of ageing effects using semi-permeable membrane devices
P. Menchai, L. Van Zwieten, S. Kimber, N. Ahmad, P.S.C. Rao, G. Hose

Bioavailable DDT assessed by SPMDs is reduced when contaminated soil is incubated as sediment.

119 Microbial and plant ecology of a long-term TNT-contaminated site
E.R. Travis, N.C. Bruce, S.J. Rosser

Long-term contamination of soil with TNT reduces the extent and diversity of vegetation, decreases culturable bacterial numbers and shifts the microbial community composition.

127 Specific accumulation of arsenic compounds in green turtles (*Chelonia mydas*) and hawksbill turtles (*Eretmochelys imbricata*) from Ishigaki Island, Japan
T. Agusa, K. Takagi, R. Kubota, Y. Anan, H. Iwata, S. Tanabe

Green turtles and hawksbill turtles have specific accumulation features of arsenic.

137 Arbuscular mycorrhizal fungi differentially affect the response to high zinc concentrations of two registered poplar clones
G. Lingua, C. Franchin, V. Todeschini, S. Castiglione, S. Biondi, B. Burlando, V. Parravicini, P. Torrigiani, G. Berta

Inoculation with arbuscular mycorrhizal fungi can improve poplar tolerance to heavy metals in phytoremediation programmes.

148 Influence of organic amendments on diuron leaching through an acidic and a calcareous vineyard soil using undisturbed lysimeters
M. Thevenot, S. Dousset, S. Rousseaux, F. Andreux

The application of organic amendments increased diuron leaching through a sandy–loam soil, in contrast to a clay–loam soil.

157 Assessment of biological effects of environmental pollution along the NW Mediterranean Sea using red mullets as sentinel organisms
I. Zorita, M. Ortiz-Zarragoitia, I. Apraiz, I. Cancio, A. Orbea, M. Soto, I. Marigómez, M.P. Cajaraville

Application of biomarkers in red mullets is promising to assess environmental pollution in the NW Mediterranean Sea.

169 Monitoring contaminants from oil production at sea by measuring gill EROD activity in Atlantic cod (*Gadus morhua*)
A. Abrahamson, I. Brandt, B. Brunström, R.C. Sundt, E.H. Jørgensen

Gill EROD activity is a sensitive biomarker for CYP1A-inducing pollutants in cod.

176 Heavy metal concentrations in marine molluscs from the Moroccan coastal region
M. Maanan

The accumulated metal concentrations in bivalves from the clean stations may be considered as useful Atlantic coast reference background levels for future comparison.

184 Bioaccumulation and toxicity of a cationic surfactant (DODMAC) in sediment dwelling freshwater invertebrates
S.D.W. Comber, K.L. Rule, A.U. Conrad, S. Höss, S.F. Webb, S. Marshall

The cationic surfactant, DODMAC, exhibits low bioavailability and toxicity to sediment dwelling organisms, with uptake dominated by ingestion.

192 AHH activity, tissue dose and DNA damage in different tissues of the scallop *Chlamys farreri* exposed to benzo[*a*]pyrene
L. Pan, J. Miao, J. Wang, J. Liu

The combination of AHH activity, B[a]P accumulation and DNA damage in the scallop Chlamys farreri can be a reasonable biomarker battery for pollution monitoring.

199 Sensitivity of submersed freshwater macrophytes and endpoints in laboratory toxicity tests
G.H.P. Arts, J.D.M. Belgers, C.H. Hoekzema, J.T.N.M. Thissen

A range of endpoints is more representative of macrophyte fitness than biomass and growth only.

207 The impact of synthetic pyrethroid and organophosphate sheep dip formulations on microbial activity in soil
T.K. Boucard, C. McNeill, R.D. Bardgett, C.D. Paynter, K.T. Semple

The application of sheep dip formulations can have a profound impact upon microbial activity and substrate utilisation in soil.

215 Seasonal and particle size-dependent variations in gas/particle partitioning of PCDD/Fs
S.-J. Lee, D. Ale, Y.-S. Chang, J.-E. Oh, S.K. Shin

Gas/particle partitioning of atmospheric PCDD/Fs for different particle sizes reflects the impacts of emitters of different size ranges.

223 Spatial variations and temporal trends between 1994 and 2005 in polychlorinated biphenyls, organochlorine pesticides and heavy metals in European eel (*Anguilla anguilla* L.) in Flanders, Belgium
J. Maes, C. Belpaire, G. Goemans

Although tissue concentrations of PCBs, pesticides and heavy metals decreased over time, they remain sufficiently high for the consumption of wild captured eel to still be discouraged.

238 Activity-dependent labeling of oxygenase enzymes in a trichloroethene-contaminated groundwater site
M.H. Lee, S.C. Clingenpeel, O.P. Leiser, R.A. Wymore, K.S. Sorenson Jr., M.E. Watwood

Enzyme probes provide a direct means of assessing aerobic cometabolic activity and a tool that enhances natural attenuation evaluations, an area lacking advancement for years.

247 Assessment of population exposure to particulate matter pollution in Chongqing, China
S. Wang, Y. Zhao, G. Chen, F. Wang, K. Aunan, J. Hao

Using an indirect microenvironment model, the population weighted exposure (PWE) to PM_{10} in Chongqing was estimated to be 211 $\mu g/m^3$ with significant contribution from indoor pollution.

Number 2 2008

Regular papers

257 Distribution of total and methylmercury in different ecosystem compartments in the Everglades: Implications for mercury bioaccumulation
G. Liu, Y. Cai, T. Philippi, P. Kalla, D. Scheidt, J. Richards, L. Scinto, C. Appleby

Mercury bioaccumulation in Florida Everglades is related to the distribution patterns of mercury species among ecosystem compartments.

266 Lead and polycyclic aromatic hydrocarbons (PAHs) in surface soil from day care centres in the city of Bergen, Norway
T. Haugland, R.T. Ottesen, T. Volden

Surface soil in 45% of the studied day care centres was contaminated by lead and PAH.

273 Geographical and pedological drivers of distribution and risks to soil fauna of seven metals (Cd, Cu, Cr, Ni, Pb, V and Zn) in British soils
D.J. Spurgeon, P. Rowland, G. Ainsworth, P. Rothery, S. Long, H.I.J. Black

Metal distributions and risks explained by balance of sources and soil property effects on fate.

284 The utility of *Pinus sylvestris* L. in dendrochemical investigations: Pollution impact of lead mining and smelting in Darley Dale, Derbyshire, UK
J.G.A. Lageard, J.A. Howell, J.J. Rothwell, I.B. Drew

Mean site dendrochemical records for Pinus sylvestris can be used to estimate the general scale and timing of atmospheric pollution episodes.

295 Effect of vegetation type on throughfall deposition and seepage flux
A. De Schrijver, J. Staelens, K. Wuyts, G. Van Hoydonck, N. Janssen, J. Mertens, L. Gielis, G. Geudens, L. Augusto, K. Verheyen

Vegetation type determines the time of onset of nitrogen saturation.

304 Metal homeostasis in *Hypogymnia physodes* is controlled by lichen substances
M. Hauck

Lichen substances in Hypogymnia physodes specifically reduce the uptake of Cu^{2+} and Mn^{2+}, but not Fe^{2+} or Zn^{2+}.

309 Genotypic and environmental variation in chromium, cadmium and lead concentrations in rice
F. Zeng, Y. Mao, W. Cheng, F. Wu, G. Zhang

Some rice genotypes had consistently low grain Cr, Cd and Pb concentrations under the soil with differently contaminated levels.

315 Alterations in soil microbial activity and *N*-transformation processes due to sulfadiazine loads in pig-manure
A. Kotzerke, S. Sharma, K. Schauss, H. Heuer, S. Thiele-Bruhn, K. Smalla, B.-M. Wilke, M. Schloter

This study gives first insights into the effects of manure containing the antibiotic sulfadiazine on microbial activity and nitrogen transformation potentials in soil.

323 Uptake and localisation of lead in the root system of *Brassica juncea*
D.E.R. Meyers, G.J. Auchterlonie, R.I. Webb, B. Wood

The sites of Pb sequestration within the root system of hydroponically grown Brassica juncea were identified.

333 Role of Chinese wind-blown dust in enhancing environmental pollution in Metropolitan Seoul
W. Kim, S.-J. Doh, Y. Yu, M. Lee

Magnetic quantification of anthropogenic pollution of Asian dust.

342 Effects of modern and ancient human activities on mercury in the environment in Xi'an area, Shannxi Province, P.R. China
Y. Jin, X. Wang, J. Lu, C. Zhang, Q. Duan

Urban activities are sources of mercury to the environment and the pomegranates grown over the burial mound of the Qinshihuang Mausoleum are not mercury-contaminated.

351 Soil metal concentrations and vegetative assemblage structure in an urban brownfield
F.J. Gallagher, I. Pechmann, J.D. Bogden, J. Grabosky, P. Weis

The study yields insight into the impact of metal contaminates soils on vegetative assemblage structure and development.

362 Screening the phytoremediation potential of desert broom (*Baccharis sarothroides* Gray) growing on mine tailings in Arizona, USA
N. Haque, J.R. Peralta-Videa, G.L. Jones, T.E. Gill, J.L. Gardea-Torresdey

Desert broom, a potential hyperaccumulating plant to clean up Cu, Pb, Cr, Zn, As, Ni and Co from the mine tailings in AZ, USA.

369 Effects of copper on spore germination, growth and ultrastructure of *Polypodium cambricum* L. gametophytes
S. Muccifora

Copper excess induced changes in development of Polypodium cambricum gametophytes, possibly by re-orientation of growth.

376 Leaf accumulation of trace elements and polycyclic aromatic hydrocarbons (PAHs) in *Quercus ilex* L.
F. De Nicola, G. Maisto, M.V. Prati, A. Alfani

Different Quercus ilex leaf accumulation of trace elements and PAHs.

384 Occurrence and fate of alkylphenols and alkylphenol ethoxylates in sewage treatment plants and impact on receiving waters along the Ter River (Catalonia, NE Spain)
R. Céspedes, S. Lacorte, A. Ginebreda, D. Barceló

Whereas partitioning of nonylphenol upon particulate matter and enrichment in sludge has been detected, long chain NPE_nOs were mainly found in the dissolved phase.

393 Application of biochemical fingerprinting and fatty acid methyl ester profiling to assess the effect of the pesticide Atradex on aquatic microbial communities
J.G. Littlefield-Wyer, P. Brooks, M. Katouli

Atradex at concentration above 245 mg L^{-1} may irreversibly change the structure and functional status of aquatic microbial communities.

401 Bioaccumulation of persistent organic pollutants in female common dolphins (*Delphinus delphis*) and harbour porpoises (*Phocoena phocoena*) from western European seas: Geographical trends, causal factors and effects on reproduction and mortality
G.J. Pierce, M.B. Santos, S. Murphy, J.A. Learmonth, A.F. Zuur, E. Rogan, P. Bustamante, F. Caurant, V. Lahaye, V. Ridoux, B.N. Zegers, A. Mets, M. Addink, C. Smeenk, T. Jauniaux, R.J. Law, W. Dabin, A. López, J.M. Alonso Farré, A.F. González, A. Guerra, M. García-Hartmann, R.J. Reid, C.F. Moffat, C. Lockyer, J.P. Boon

High PCB levels were recorded in porpoises and common dolphins from European coasts.

416 Multiple experimental approaches of immunotoxic effects of mercury chloride in the blue mussel, *Mytilus edulis*, through in vivo, in tubo and in vitro exposures
M.B. Duchemin, M. Auffret, N. Wessel, M. Fortier, Y. Morin, J. Pellerin, M. Fournier

$HgCl_2$ impairs the immune system of test-tube mussels.

424 Bioaccumulation of polychlorinated biphenyls in the eel (*Anguilla anguilla*) at the Camargue Nature Reserve – France
C.A. Oliveira Ribeiro, Y. Vollaire, E. Coulet, H. Roche

The reserve of Camargue – South of France is impacted by a myriad of pollutant organic persistent like PCBs.

432 Short term exposure to elevated trinitrotoluene concentrations induced structural and functional changes in the soil bacterial community
E.R. Travis, N.C. Bruce, S.J. Rosser

The bacterial community structure, function and diversity changed when soil was amended with TNT; pseudomonads increased in prevalence and tolerance to TNT toxicity increased.

440 Evaluation of stream ecological integrity using litter decomposition and benthic invertebrates
J. Castela, V. Ferreira, M.A.S. Graça

This study supports the need for incorporating functional measures in evaluations of stream ecological integrity.

450 Effects of three pesticides on the avoidance behavior of earthworms in laboratory tests performed under temperate and tropical conditions
M. Garcia, J. Römbke, M.T. de Brito, A. Scheffczyk

The earthworm avoidance test is a practical and sensitive screening method for assessing the effects of pesticides in tropical soils.

457 Vertical distribution and environmental significance of sulfur and oxygen heterocyclic aromatic hydrocarbons in soil samples collected from Beijing, China
Z. Zhihuan, H. Fengpeng, B. Qingwei, L. Song

The vertical distribution of SOHAHs was provided and possible sources were different between topsoils and deep part.

468 Bioavailability assessment of contaminants in soils via respiration and nitrification tests
K. Hund-Rinke, M. Simon

The paper provides a further criterion for a more detailed assessment of soil contamination, leading to different recommended actions depending on which parameter is affected.

476 Heavy metals' content of automotive shredder residues (ASR): Evaluation of environmental risk
O. Gonzalez-Fernandez, M. Hidalgo, E. Margui, M.L. Carvalho, I. Queralt

High amounts of Pb and Zn from the finest fractions of automotive shredder residue imply environmental and toxic risks.

483 Removal of phenolic estrogen pollutants from different sources of water using molecularly imprinted polymeric microspheres
Y. Lin, Y. Shi, M. Jiang, Y. Jin, Y. Peng, B. Lu, K. Dai

Bisphenol A imprinted polymeric microspheres were used as absorbent and could remove phenolic estrogen pollutants from water effectively.

Short communication

492 Flow cytometric measurement of the clearance rate in the blue mussel *Mytilus edulis* and the development of a new individual exposure system for aquatic immunotoxicological studies
M.B. Duchemin, N. Wessel, M. Fournier, M. Auffret

Test-tube mussels offer a new exposure design for immunotoxicology.

Number 3 2008

Review

497 Performance of bioaugmentation-assisted phytoextraction applied to metal contaminated soils: A review
T. Lebeau, A. Braud, K. Jézéquel

Bioaugmentation-assisted plant improves the phytoextraction performances for soils contaminated by metals.

New initiatives

523 **Low toxicant concentrations decrease the frequency of fast-growing seedlings at high densities of annual baby's breath (*Gypsophila elegans*)**
A. Sinkkonen, R. Strömmer, O.-P. Penttinen

Very low toxicant concentrations reduce the frequency of most vital seedlings in dense stands in vitro.

526 **Deriving ozone dose–response of photosynthesis in adult forest trees from branch-level cuvette gas exchange assessment**
C. Then, M. Löw, R. Matyssek, G. Wieser

Branch-level O_3 dose dependence of photosynthesis derived from cuvette assessment yields sun-crown foliage sensitivity under whole-tree free-air O_3 fumigation.

Regular papers

529 **Snapping turtles (*Chelydra serpentina*) as bioindicators in Canadian Areas of Concern in the Great Lakes Basin. II. Changes in hatching success and hatchling deformities in relation to persistent organic pollutants**
S.R. de Solla, K.J. Fernie, S. Ashpole

Hatching success and deformities of snapping turtle eggs varied among Great Lake Areas of Concern, but were not attributable to specific chemical exposure.

537 **Long-term Hg pollution induced Hg tolerance in the terrestrial isopod *Porcellio scaber* (Isopoda, Crustacea)**
A. Lapanje, D. Drobne, N. Nolde, J. Valant, B. Muscet, V. Leser, M. Rupnik

Isopods (Porcellio scaber) as well as their bacterial gut community from a mercury-polluted site are mercury tolerant.

548 **Prediction of Pb speciation in concentrated and dilute nutrient solutions**
P.M. Kopittke, C.J. Asher, N.W. Menzies

Due to the low solubility of Pb-phosphates, nutrient solutions with low P concentrations should be used for investigating the phytotoxicity of Pb.

555 **Effect of citric acid and EDTA on chromium and nickel uptake and translocation by *Datura innoxia***
L. Jean, F. Bordas, C. Gautier-Moussard, P. Vernay, A. Hitmi, J.-C. Bollinger

Chelant effectiveness in increasing chromium and nickel uptake by Datura innoxia is the result of the increase in translocation versus negative effect on plant physiology.

564 **Growth responses of low-alpine dwarf-shrub heath species to nitrogen deposition and management**
A.J. Britton, J.M. Fisher

Low levels of N deposition increase productivity in alpine dwarf-shrub heath despite strong climatic constraints.

574 **ELISA detection of multixenobiotic resistance transporter induction in indigenous freshwater Chironomidae larvae (Diptera): A biomarker calibration step for *in situ* monitoring of xenobiotic exposure**
X. Moreau, G. Saez, A. Thiéry, O. Clot-Faybesse, G. Guiraudie-Capraz, C. Bienboire-Frosini, C. Martin, L. De Jong

Semi-quantification of midge larval MXR transporters by ELISA is a simple and sensitive method to detect chemically polluted situations in running freshwaters.

582 **Peat soils as a source of lead contamination to upland fluvial systems**
J.J. Rothwell, M.G. Evans, S.M. Daniels, T.E.H. Allott

Erosion can transform peat soils from sinks to sources of lead contamination.

590 **Genotoxicity detected in wild mice living in a highly polluted wetland area in south western Spain**
S. Mateos, P. Daza, I. Domínguez, J.A. Cárdenas, F. Cortés

We have found an increased genotoxic damage in wild mice in a highly polluted area from industry, mining and agriculture in SW Spain, as assessed by the Comet assay.

594 **Source diagnostics of polycyclic aromatic hydrocarbons in urban road runoff, dust, rain and canopy throughfall**
W. Zhang, S. Zhang, C. Wan, D. Yue, Y. Ye, X. Wang

Urban road runoff and road dust, canopy throughfall and rain were considered as a system for diagnostics of PAH sources.

602 **Physiological responses to atmospheric fluorine pollution in transplants of *Pseudoscleropodium purum***
J.R. Aboal, J.A. Couto, J.A. Fernández, A. Carballeira

Good correlations between fluorine concentrations in transplants of terrestrial moss and chlorophyll contents and pigment stress indexes were observed.

610 **Levels and congener profiles of polybrominated diphenyl ethers (PBDEs) in Zebra mussels (*D. polymorpha*) from Lake Maggiore (Italy)**
A. Binelli, L. Guzzella, C. Roscioli

This study shows the first data of PBDE contamination in freshwater invertebrates from Mediterranean basin.

618 **Geographic distribution of selected elements in the livers of polar bears from Greenland, Canada and the United States**
S.A. Rush, K. Borgå, R. Dietz, E.W. Born, C. Sonne, T. Evans, D.C.G. Muir, R.J. Letcher, R.J. Norstrom, A.T. Fisk

Geographical trends were observed for a number of elements in livers, including mercury, of polar bears collected across Alaska, Canada and Greenland and were similar to those observed in the early 1980s.

627 **A model compound study: The ecotoxicological evaluation of five organic contaminants employing a battery of marine bioassays**
A. Macken, M. Giltrap, B. Foley, E. McGovern, B. McHugh, M. Davoren

Ecotoxicological evaluation of five organic marine sediment contaminants was conducted and the suitability of the test species for marine porewater TIE discussed.

638 **Immunological alterations in juvenile Pacific herring, *Clupea pallasi*, exposed to aqueous hydrocarbons derived from crude oil**
C.J. Kennedy, A.P. Farrell

Effects of oil on herring immunology and disease resistance.

649 **Field monitoring of volatile organic compounds using passive air samplers in an industrial city in Japan**
K. Kume, T. Ohura, T. Amagai, M. Fusaya

Passive air samplings with hood are useful for determining the identities, sources, and distributions of ambient VOC pollutants.

658 **Effect of two major *N*-nitroso hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) metabolites on earthworm reproductive success**
B. Zhang, S.B. Cox, S.T. McMurry, W.A. Jackson, G.P. Cobb, T.A. Anderson

RDX metabolites affect earthworm cocoon production and hatching.

668 **Plant uptake of trace elements on a Swiss military shooting range: Uptake pathways and land management implications**
B.H. Robinson, S. Bischofberger, A. Stoll, D. Schroer, G. Furrer, S. Roulier, A. Gruenwald, W. Attinger, R. Schulin

Establishment of a complete vegetation cover on shooting ranges would reduce the contamination of plant leaves by toxic trace elements.

677 **Environmental risks of farmed and barren alkaline coal ash landfills in Tuzla, Bosnia and Herzegovina**
A. Dellantonio, W.J. Fitz, H. Custovic, F. Repmann, B.U. Schneider, H. Grünewald, V. Gruber, Z. Zgorelec, N. Zerem, C. Carter, M. Markovic, M. Puschenreiter, W.W. Wenzel

Uncontrolled farming and tillage of previously soil-covered coal ash landfills resulted in exposure of ash on the surface.

687 **Oxidative stress in duckweed (*Lemna minor* L.) caused by short-term cadmium exposure**
J. Razinger, M. Dermastia, J.D. Koce, A. Zrimec

The increase of total antioxidative potential at low cadmium concentration is one of the mechanisms that helps duckweed to cope with cadmium-induced oxidative stress.

Short communications

695 **Genotoxicity evaluation of the insecticide endosulfan in the wetland macrophyte *Bidens laevis* L.**
D.J. Pérez, M.L. Menone, E.L. Camadro, V.J. Moreno

Endosulfan causes a concentration-dependent increase of chromosome aberrations in the macrophyte Bidens laevis.

699 **Spore release by the green alga *Ulva*: A quantitative assay to evaluate aquatic toxicants**
T. Han, Y.-S. Han, C.Y. Park, Y.S. Jun, M.J. Kwon, S.-H. Kang, M.T. Brown

A simple and cost-effective bioassay using spore release by the green macroalga Ulva has been developed and the sensitivity is similar to or greater than other well-established tests.

706 **Effect of housing geometry on the performance of Chemcatcher™ passive sampler for the monitoring of hydrophobic organic pollutants in water**
T. Lobpreis, B. Vrana, E. Dominiak, K. Dercová, G.A. Mills, R. Greenwood

The effect of passive sampler geometry on accumulation kinetics of organic pollutants from water was evaluated.

Corrigendum

711 **Corrigendum to "Estimates of ambient background concentrations of trace metals in soils for risk assessment" [Environ Pollut 148 (2007) 221–229]**
F.J. Zhao, S.P. McGrath, G. Merrington

I **Contents of Volume 153**